

Development of Kolmogorov spectrum in the pulsar radio emission

G. Z. Machabeli and G. Gogoberidze*

*Center for Plasma Astrophysics, Abastumani Astrophysical Observatory,
Ave. A. Kazbegi 2, Tbilisi 0160, Georgia*

(Dated: February 2, 2008)

Abstract

It is shown that the scattering of electromagnetic waves by Langmuir ones taking into the account the electric drift motion of particles, is the most intense nonlinear process and should be responsible for the formation of radiation spectrum of radio pulsars. Performed analysis indicates that in the case of existence of inertial interval formation of stationary spectra is possible. Analysis of linear mechanisms of wave generation and silk allow to conclude that only possible stationary solution has spectral index $\alpha = -1.5$. Obtained spectrum is in good agreement with observational data.

PACS numbers: 97.60.Gb, 52.35.Ra

*Electronic address: gogober@geo.net.ge

I. INTRODUCTION

Aim of the present work is to study nonlinear processes responsible for the formation of the radio emission spectrum in the pulsar magnetosphere. The spectra of pulsar radio emission has been intensively studied by observers [1, 2, 3, 4, 5]. Above 100 MHz observed intensity I of the majority of pulsars can be described by a simple power law $I \sim \nu^\alpha$ with average value of spectral index $\langle\alpha\rangle = -1.8 \pm 0.2$ [2]. Spectral index of individual pulsars vary from $\alpha = +1.4$ to $\alpha = -3.8$. Some pulsars has the spectra which can be modelled by the two power law [1, 3], but it seems that single power law is a rule and the two power law spectrum is a rather rare exception [2].

From our point of view pulsar radio emission is generated and formed in the magnetosphere. Therefore properties of the radiation is mainly determined by the physical processes in the magnetosphere plasma. This approach allows to explain other main characteristics of pulsar radiation, such as polarization properties [6], nullings [7], micro impulses [8], mode switching [9] and so on.

In the presented paper we study nonlinear processes in the magnetosphere plasma that should be responsible for the formation of the pulsar radio emission spectra. Mathematical methods used for analysis is based on the weak turbulence theory (see, e.g., [10, 11]).

The paper is organized as follows: the model of the magnetosphere as well as linear eigenmodes of e^-e^+ plasma and linear mechanisms of their generation is discussed in Sec. 2. Nonlinear processes are studied in Sec. 3. In Sec 4 stationary solutions for the spectra are obtained and analyzed.

II. LINEAR MODES AND THEIR GENERATION IN PULSAR MAGNETOSPHERE

According to the standard model of the pulsar magnetosphere [12, 13] relativistic e^-e^+ plasma that is pierced by electron beam moves along open field lines of pulsar magnetic field $\mathbf{B}_0$. Magnetic field of pulsar is supposed to be dipole. Under open field lines we mean the field lines that crosses the surface of light cylinder - the surface where the velocity of field line solid rotation reaches the speed of light c . It is supposed that electric field directed along the open field lines is generated at the surface of pulsar [12], that pulls out electrons of the

primary beam from the surface of the star. The particles moving along magnetic field lines generate γ rays, which on its turn if the energy $\epsilon_\gamma > 2m_e c^2$ generate electron-positron pairs [13]. This process is stopped when the initial electric field becomes screened. The particle distribution function that has the form presented in Fig. 1 can be presented as:

$$f = f_p + f_t + f_b, \quad (1)$$

where f_p describes the bulk of the plasma, f_t is the distribution function of elongated to the plasma motion tail particles and f_b corresponds to the particles of the primary beam. The distribution function (1) is one dimensional. In Fig. 1 solid and dashed lines correspond to electrons and positrons respectively. The shift of the distribution functions is caused by the existence of the primary beam. It is supposed that

$$n_p \gamma_p \approx n_t \gamma_t \approx \frac{1}{2} n_b \gamma_b, \quad (2)$$

where $\gamma_p, \gamma_t, \gamma_b$ are Lorentz factors and n_p, n_t, n_b concentrations of bulk, tail and beam particles respectively. For typical pulsars $\gamma_b \sim 10^6 - 10^7$, $n_b^0 \sim 10^{11}$, $\gamma_t \sim 10^4$, $n_t^0 \sim 10^{13} - 10^{14}$, $\gamma_p \sim 10$, $n_p^0 \sim 10^{17}$. Index 0 indicates that the values belong to the surface of the pulsar.

The condition of quasi neutrality yields

$$\Delta\gamma = \gamma_+ - \gamma_- = \int f_+ \gamma d^3p - \int f_- \gamma d^3p, \quad (3)$$

here indexes $\pm$ are related to positrons and electrons respectively. $f_\pm$ are normalized such that $\int f_\pm d^3p = 1$.

In spite of the fact that $\Delta\gamma \ll \gamma_\pm$ this small difference plays important role in explanation of polarization properties of pulsar radiation. The radiation of the most pulsars has considerable part of circularly polarized radiation. It was shown [6] that circular polarization can occur only in small angle θ between wave vector $\mathbf{k}$ and $\mathbf{B}_0$:

$$\theta^2 \lesssim \frac{\omega}{\omega_B} \Delta\gamma \ll 1. \quad (4)$$

Consequently, the observer receives the radiation that is formed in small angle with respect to $\mathbf{B}_0$. So, for pulsar radio emission the waves propagating nearly along the magnetic field are important. This agrees with linear theory of e^-e^+ plasma in strong magnetic field: analysis provides that the wave generation takes place only almost along the magnetic field [6, 14].

Due to the absence of gyrotropy the spectra of linear modes in e^-e^+ plasma is meager. There exist only three modes [14]: purely transversal t mode and in general partially potential lt modes. Electric vector of t mode is perpendicular to both $\mathbf{k}$ and $\mathbf{B}_0$. If $\mathbf{k} \parallel \mathbf{B}_0$ the dispersion relation of this mode is:

$$\omega_t = kc(1 - \delta), \quad (5)$$

where $\delta \equiv \omega_p^2/(4\omega_B^2\langle\gamma^3\rangle)$, and ω_p , ω_B are plasma and cyclotron frequencies respectively

$$\omega_p \equiv \left(\frac{8\pi e^2 n_p}{m_e}\right)^{1/2}, \quad \omega_B \equiv \frac{e|B_0|}{m_e c}, \quad (6)$$

angular brackets denote averaging over distribution function and $\langle\gamma\rangle \approx \gamma_p$.

lt mode has two branches lt_1 and lt_2 . Electric vectors of these modes are located in the plane formed by $\mathbf{k}$ and $\mathbf{B}_0$. If $\mathbf{k} \parallel \mathbf{B}_0$ mode lt_2 merges with t mode with spectrum (5). The mode lt_1 propagating along $\mathbf{B}_0$ is longitudinal Langmuir wave (l). If $\omega \gtrsim kc$ their dispersion is [15, 16]:

$$\omega_l^2 = \omega_p^2\langle\gamma^{-3}\rangle + 3k^2c^2 \left(1 - \frac{\langle\gamma^{-5}\rangle}{\langle\gamma^{-3}\rangle}\right), \quad (7)$$

and when $\omega \approx kc$:

$$\omega_l \approx c[k - \alpha(k - k_0)], \quad (8)$$

where $k_0^2 = 2\langle\gamma\rangle\omega_p^2/c^2$ and $\alpha = \langle\gamma\rangle/2\langle\gamma^3\rangle$.

From Eqs. (5) and (8) it follows that frequencies of l and t modes become equal at:

$$k \approx \bar{k} \equiv k_0 \left(1 + \frac{1}{2} \frac{\omega_p^2}{\omega_B^2} \langle\gamma^3\rangle\right), \quad (9)$$

Dispersion relations of the waves propagating along $\mathbf{B}_0$ are presented in Fig. 2.

Mechanisms of wave generation in the magnetosphere e^-e^+ plasma has been intensively studied (see, e.g., [17]). The magnetosphere plasma can be unstable due to two reasons: a) existence of energetic beam; b) absence of symmetry and one dimensionality of the distribution function. Detailed analysis provides [6] that the beam instability can develop only due to the drift of the particles in the curved magnetic field (so called "drift-cherenkov" resonance). Linear stage of this instability was studied in [19] and nonlinear stage in [20]. In this paper we will be interested in the second mechanism of instability - excitation at the anomalous Doppler-effect resonance. The condition of this resonance is [6]:

$$\omega - k_{\parallel}v_{\parallel} - k_{\perp}u_{\perp}^m + \frac{\omega_B}{\gamma_r} \approx 0. \quad (10)$$

Here $k_{\parallel}$ and $k_{\perp}$ are projections of wave vector along and perpendicular to $\mathbf{B}_0$, $u_{\perp}^m \equiv cv_{\parallel}\gamma_r/\omega_B R_B$ is curvature drift velocity, R_B is curvature radius of magnetic field lines and γ_r is Lorentz factor of resonant particles.

Detailed analysis provides that this condition can be satisfied only for energetic particles of the tail and beam [21], when the condition $2\gamma_r^2\delta > 1$ is satisfied. Substituting Eq. (2) into (10), for $k_{\perp} = 0$ we obtain:

$$\omega_t \approx k_{\parallel}c \approx \frac{\omega_B}{\gamma_r\delta}. \quad (11)$$

From this equation for typical pulsar parameters [22, 23] it follows that the frequencies of generated t waves

$$\omega_t \ll \omega_B. \quad (12)$$

It should be noted that transversal waves are strongly damped by cyclotron damping with the bulk particles. The condition of this resonance yields

$$\omega - k_{\parallel}v_{\parallel} - k_{\perp}u_{\perp}^m - \frac{\omega_B}{\gamma_r} \approx 0. \quad (13)$$

This condition provides that cyclotron damping is effective for the waves with frequencies

$$\omega_t \gtrsim 2\gamma_p\omega_B. \quad (14)$$

Linear mechanisms of the wave generation discussed in this section emerges transversal t waves. On its turn, nonlinear interactions of waves redistributes the energy among different modes and scales and as it will be shown bellow can lead to the formation of stationary spectra of wave turbulence.

III. NONLINEAR PROCESSES

For the development of weak turbulence theory the small parameter of weak turbulence is used. In the case of e^-e^+ plasma this parameter is

$$\frac{|E|^2}{m_e c^2 n_p \gamma_p} \ll 1, \quad (15)$$

where $\mathbf{E}$ is electric vector of the wave. This condition implies that we consider the state when the energy density of plasma is much greater then the energy density of waves. except the parameter (15) in the magnetosphere plasma there exist another two small parameters

that greatly simplify future analysis. The first small parameter is $\omega_p^2/\omega_B^2 \ll 1$. The second one:

$$\gamma_p^2 \frac{(\omega - k_{\parallel} v_{\parallel})^2}{\omega_B^2} \ll 1 \quad (16)$$

allows us to use so-called drift approximation [24].

As it is well known [25] if turbulence is weak the dynamics is governed by three wave resonant processes. In e^-e^+ plasma there exists additional limitations for realization of three wave resonant interactions. The absence of gyrotopry causes the fact that the terms proportional to the odd power of the charge do not contribute to the nonlinear currents. Another specific property of the magnetosphere plasma is that all the waves are propagating almost along the magnetic field.

Breakup of l wave into two t waves was studied in [26], and all possible three wave processes in [27]. In this researches main attention was payed to possible breakup of l waves because it was believed that these waves should be the most unstable ones. Existence of the anomalous Doppler-effect resonance discussed in the previous Section leads to the conclusion that in fact the most interesting three wave processes are possible breakups of t waves that should be responsible for the formation of observed spectra of pulsars.

For the analysis of nonlinear processes it is useful to introduce quantities and notations used in quantum physics. Usually plasmon occupation numbers N_k^λ and wave amplitudes a_k^λ are introduced [11, 24, 25]:

$$\omega_k N_k^\lambda \equiv \omega_k \langle a_k^\lambda a_k^{\lambda*} \rangle = \frac{1}{\omega_k} \left(\frac{\partial}{\partial \omega} \omega^2 \varepsilon_k^\lambda \right)_{\omega=\omega_k^\lambda} \frac{|E_k|^2}{4\pi}. \quad (17)$$

here

$$\varepsilon_k^\lambda \equiv \frac{e_i^{\lambda*} e_j^\lambda}{\omega^2} (k_i k_j c^2 + \omega^2 \varepsilon_{ij}); \quad (18)$$

$\mathbf{e}^\lambda$ is polarization vector of $\lambda(lt, l, t)$ mode; $E_j^\lambda \equiv e_j^\lambda E$; ε_{ij} is dielectric tensor; $*$ denotes complex conjugation and angular brackets denote averaging over phases.

Maxwell equations yield

$$\begin{aligned} \partial_t a_k + i\omega_k^\lambda a_k &= \sum_{\lambda_1 \lambda_2} \int d\mathbf{k}_1 d\mathbf{k}_2 d\omega_1 d\omega_2 a_k^{\lambda_1} a_k^{\lambda_2*} \\ &\times \delta(\mathbf{k} - \mathbf{k}_1 - \mathbf{k}_2) \delta(\omega - \omega_1 - \omega_2) V_{\lambda|\lambda_1 \lambda_2}, \end{aligned} \quad (19)$$

where matrix element of the process $\lambda \rightarrow \lambda_1 + \lambda_2$

$$\begin{aligned} V_{\lambda|\lambda_1 \lambda_2} &= 4\pi \frac{\omega_{k_1}^{\lambda_1} \omega_k^\lambda}{\omega_k^\lambda} (\omega_{k_1}^{\lambda_1} - \omega_{k_2}^{\lambda_2}) (\partial_\omega \omega^2 \varepsilon^\lambda)^{-1/2}_{\omega=\omega_k^\lambda} \\ &\times (\partial_\omega \omega^2 \varepsilon^\lambda)^{-1/2}_{\omega=\omega_{k_1}^{\lambda_1}} (\partial_\omega \omega^2 \varepsilon^\lambda)^{-1/2}_{\omega=\omega_{k_2}^{\lambda_2}} \sigma_{\lambda|\lambda_1 \lambda_2}, \end{aligned} \quad (20)$$

and $\sigma_{\lambda|\lambda_1\lambda_2}$ is nonlinear conductivity tensor.

If $k_\perp = 0$, the process $t \rightarrow t' + t''$ is forbidden due to the fact that the second order current $j^{(2)} \sim e^3$. When $k_\perp \neq 0$, this process can take place but the matrix element of interaction is negligibly small [27]: $V_{t|t't''} \sim \omega_p^3/\omega_B^3 \ll 1$.

Another possible three wave process $t \rightarrow t' + l$ was discussed in [27] without taking into consideration drift motion of the particles. In this case the frequency of generated Langmuir waves $\omega_l \approx k_l c$. If all the waves are propagating along $\mathbf{B}_0$, matrix element of interaction is

$$V_{t|lt'} \approx \frac{e}{m_e c} \frac{\omega_p^2}{\omega_B \omega_l} \sqrt{\frac{\omega_t \omega_{t'}}{\omega_l}}. \quad (21)$$

Consideration of the drift motion of particles makes this process even more intense. For simplicity let us consider the case when the waves are propagating along $\mathbf{B}_0$. Let us assume that electric vectors of t and t' waves are directed along x and y axes respectively (see Fig. 3) and $\mathbf{B}_0$ is parallel to z axis. E_x^t causes the drift motion of particles along y :

$$u_y^d = c \frac{E_x^t}{B_0}, \quad (22)$$

which on its turn generates nonlinear electric field:

$$\mathbf{E} = \frac{1}{c} [\mathbf{u}^d, \mathbf{B}']. \quad (23)$$

It has to be noted that drift velocity is the same for electrons and positrons. Consequently the drift motion do not lead to current generation in the linear approximation. If the resonant conditions

$$\omega_t = \omega_l + \omega_{t'}, \quad \mathbf{k}_t = \mathbf{k}_l + \mathbf{k}_{t'}, \quad (24)$$

are fulfilled the beating of t and t' waves that generates longitudinal electric field (23) are in resonance with corresponding l wave.

Analysis of the resonant conditions (24) yields that l wave should have phase velocity a bit less then the speed of light. But this kind of l waves in general can be strongly damped by collisionless Landau damping. In this case instead of three wave nonlinear interaction one has to consider scattering of t waves by plasma particles - so called wave-particle-wave interaction [24, 25].

Analysis of standard pulsar parameters yield that in principle both cases are possible. In this paper we consider the case when l waves involved in resonance with t waves are not strongly damped by collisionless damping.

Maxwell equations governing the dynamics in the case when all the waves are propagating along $\mathbf{B}_0$ yield:

$$\left(\frac{\partial^2}{\partial t^2} - c^2 \frac{\partial^2}{\partial z^2} \right) \mathbf{E}_\perp + 4\pi \frac{\partial \mathbf{j}_\perp^{NL}}{\partial t} = 0, \quad (25)$$

$$\frac{\partial E_z}{\partial t} + 4\pi j_z^{NL} = 0. \quad (26)$$

Nonlinear currents $\mathbf{j}_\perp^{NL}$ j_z^{NL} can be readily calculated as follows: substituting (22) into Eq. (23) and taking into account Faraday law we get:

$$j_z^{NL} = -\frac{i}{4\pi} \left(\frac{k_z^t}{\omega^t} E_x E_y' + \frac{k_z^{t'}}{\omega^{t'}} E_x' E_y \right). \quad (27)$$

For $\mathbf{j}_\perp^{NL}$ we have

$$\mathbf{j}_\perp^{NL} = \mathbf{u}_\perp^d \rho_e, \quad (28)$$

where perturbation of charge density ρ_e can be determined from Poisson equation.

Substituting in Eqs. (25)-(26) electric field components as

$$E_{z,\perp}(t) = E_{z,\perp k}(t) e^{i(\omega t - k z)} + c.c., \quad (29)$$

and taking into account Eqs. (27)-(28) as well as dispersions (5) and (8) we obtain for slowly varying amplitudes:

$$\frac{\partial E_{xk_t}}{\partial t} = \frac{i}{2} \frac{c\omega_l}{B_0} \left(\frac{k^{t'}}{\omega^{t'}} + \frac{k^l}{\omega^l} \right) E_{zk_l} E_{yk_{t'}}^*, \quad (30)$$

$$\frac{\partial E_{yk_{t'}}}{\partial t} = -\frac{i}{2} \frac{c\omega_l}{B_0} \left(\frac{k^t}{\omega^t} + \frac{k^l}{\omega^l} \right) E_{zk_l} E_{xk_t}^*, \quad (31)$$

$$\frac{\partial E_{zk_l}}{\partial t} = -\frac{i}{2} \frac{c\omega_l}{\langle \gamma \rangle B_0} \left(\frac{k^t}{\omega^t} + \frac{k^{t'}}{\omega^{t'}} \right) E_{xk_t} E_{yk_{t'}}^*. \quad (32)$$

Assuming that there exist many waves with chaotic phases and introducing occupation numbers:

$$N_{x,yk} = \frac{|E_{x,yk}|^2}{4\pi\omega_t}, \quad N_{zk} = \frac{\langle \gamma^2 \rangle |E_{zk}|^2}{8\pi\omega_l}, \quad (33)$$

and using standard technique [24, 25] we get for the matrix element of interaction:

$$V_{t|t'} \approx \frac{c}{B_0} (\omega_t k_{t'} + \omega_{t'} k_t) \sqrt{\frac{\omega_l}{\omega_t \omega_{t'} \langle \gamma^2 \rangle}}. \quad (34)$$

Comparison of Eqs. (21) and (34) yields that the drift motion of particles leads to important intensification of the considered process.

IV. STATIONARY SPECTRA

In the case under consideration $\omega_t, \omega_{t'} \gg \omega_l$. Consequently, the nonlinear interaction is local - only the waves with close frequencies can effectively interact. As it was discussed in Sec. 2 the generation of t waves takes place for $\omega \ll \omega_B$, whereas the cyclotron damping is effective for $\omega > 2\gamma_p \omega_B$. Nonlinear processes are responsible for the formation of stationary spectrum between these intervals of generation and silk. As it is known [10] the stationary spectrum in the Kolmogorov interval is fully determined by the matrix element (34) and linear dispersions of waves. There exist two stationary solutions:

$$N_{k1} = A_1 k^{-s-d}, \quad N_{k2} = A_2 k^{-s-d+1/2}, \quad (35)$$

where d is the dimension of the problem and s is the index of homogeneity of the matrix element. The first solution $N_{k1} = A_1 k^{-s-d}$ corresponds to constant flux of energy to the smaller scales, whereas $N_{k2} = A_2 k^{-s-d+1/2}$ corresponds to constant flux of plasmons to the larger scales [10, 11].

In our case $d = 1$ and $S = 3/2$. Therefore $N_{k1} = A_1 k^{-5/2}$ and $N_{k2} = A_2 k^{-2}$. Taking into account the linear character of t wave dispersion for the energy spectrum $\mathcal{E}_\omega$ that is proportional to the observed intensity I_ω we have

$$I_{\omega 1} \sim \mathcal{E}_{\omega 1} \sim \omega^{-3/2}, \quad I_{\omega 2} \sim \mathcal{E}_{\omega 2} \sim \omega^{-1}. \quad (36)$$

which of this spectra is realized in practice depends on the linear mechanisms of generation and silk of the waves. In the case under consideration the waves with relatively low frequencies are excited at the anomalous doppler-effect resonance and waves with relatively high frequencies are damped at cyclotron resonance. In this case the spectrum $\mathcal{E}_{\omega 1} \sim \omega^{-3/2}$ is only possible.

As it was mentioned above most of the pulsars has the spectral indexes $-1.5 > \alpha > -2$. Consequently, obtained theoretical result $\alpha = -1.5$ seems to be in good accordance with the observations.

V. SUMMARY

Nonlinear three wave processes that can be responsible for the formation of spectra of radio pulsars is considered. Existence of strong magnetic field makes it necessary to take into

account electric drift motion of particles. In the framework of the weak turbulence theory possible stationary spectra that can occur in Kolmogorov interval are calculated. Analysis of linear mechanisms of wave generation and silk allows to conclude that only possible spectral index is $\alpha = -1.5$. Obtained result is in satisfactory accordance with the observational data.

-
- [1] V. M. Malofeev, APS Conf. Ser. **105**, 271 (1996).
 - [2] O. Maron, J. Kijak, M. Kramer and R. Wielebinski, Astron. Astrophys. **147**, 195 (2000).
 - [3] D. R. Lorimer, J. A. Yates, A. G. Lyne and M. D. Gould, Mon. Not. R. Astron. Soc. **273**, 411 (1995).
 - [4] V. A. Izvekova, A. .D. Kuzmin, V. M. Malofeev and Y. Shitov, Astrophys. and Space Science **78**, 45 (1981).
 - [5] A. G. Lyne, F. G. Smith and D. A. Graham, Mon. Not. R. Astron. Soc. **153**, 337 (1971).
 - [6] A. Z. Kazbegi, G. .Z. Machabeli and G. I. Melikidze, Mon. Not. R. Astron. Soc. **253**, 377 (1991).
 - [7] A. Kazbegi, G. Machabeli, G. Melikidze and C. Shukre, Astron. Astrophys. **309**, 515 (1996).
 - [8] G. Machabeli, D. Khechinashvili, G. Melikidze and D. Shapakidze, Mon. Not. R. Astron. Soc. **327**, 984 (2001).
 - [9] O. I. Malov, V. M. Malofeev, G. Machabeli and G. Melikidze, Astron. Zh. **74**, 303 (1997).
 - [10] V. E. Zakharov and E. A. Kuznetsov, Zh. Éksp. Teor. Fiz. **75**, 904 (1978);
 - [11] V. E. Zakharov, V.S. L'vov and G. Falkovich, *Kolmogorov Spectra of Turbulence I* (Springer-Verlag, Berlin, 1992).
 - [12] P. Goldreich and W. H. Julian, Astrphys. J. **157**, 869 (1969).
 - [13] P. A. Sturrock, Astrphys. J. **164**, 529 (1971).
 - [14] A. S. Volokitin, V. V. Krasnosel'skix and G. Z. Machabeli, Fiz. Plazmy **11**, 531 (1985).
 - [15] V. P. Silin, Zh. Éksp. Teor. Fiz. **38**, 1577 (1960).
 - [16] V. N. Tsitovich, Zh. Éksp. Teor. Fiz. **40**, 1775 (1961).
 - [17] D. B. Melrose, *Instabilities in Space and Laboratory Plasmas* (Cambridge University Press, 1986).
 - [18] V. L. Ginzburg and V. V. Zhelezniakov, Annu. Rev. Astron. Astrophys. **13**, 511 (1975).
 - [19] M. Lutikov, G. Machabeli and R. Blandford, Astrophys. J., **512**, 804 (1999).

- [20] D. Shapakidze, G. Machabeli, G. Melikidze and D. Khechinashvili, Phys. Rev. E, **67**, 345 (2003).
- [21] J. G. Lominadze, A. B. Mikhailovski and G. Z. Machabeli, Fiz. Plazmy **5**, 748 (1979).
- [22] G. Z. Machabeli and V. V. Usov, Astron. Zh. Lett. **5**, 238 (1979).
- [23] G. Z. Machabeli and V. V. Usov, Astron. Zh. Lett. **15**, 393 (1989).
- [24] V. N. Tsitovich, *Nonlinear Effects in Plasma* (Plenum, New Yourk, 1970).
- [25] R. Z. Sagdeev and A. A. Galeev, *Nonlinear Plasma Theory* (Benjamin, New Yourk, 1969).
- [26] A. B. Mikhailovski, Fiz. Plazmy **6**, 613 (1980).
- [27] M. E. Gedalin and G. Z. Machabeli, Fiz. Plazmy **9**, 1015 (1983).




FIG. 1: Particle distribution function f of magnetosphere plasma. Solid and dashed lines correspond to electrons and positrons respectively.




FIG. 2: Dispersion curves of the waves propagating along magnetic field in magnetized e^-e^+ plasma.




FIG. 3: Calculation of nonlinear currents